

First record of *Morchella pulchella* from Pakistan

H. BADSHAH¹, B. ALI¹, S.A. SHAH¹,
M.M. ALAM¹, H.I. ALY², A.S. MUMTAZ^{1*}

¹ Department of Plant Sciences, Faculty of Biological Sciences, Quaid-i-Azam University,
Islamabad, 45320, Pakistan

² Irma Lerma Rangel College of Pharmacy, Department of Pharmaceutical Sciences,
Texas A&M Health Science Center, Kingsville, TX 78363, USA

* CORRESPONDENCE TO: asmumtaz@qau.edu.pk

ABSTRACT—A species of true morel (*Morchella*) was collected in the Malam Jabba valley in the Swat District of Pakistan in April 2015. The specimen was identified by sequencing portions of RNA polymerase II largest subunit (RPB1), second largest subunit (RPB2), and translation elongation factor-1 α (TEF1). Phylogenetic analysis of the partial TEF1 sequence indicated the collection was *M. pulchella*, previously reported from China, Turkey, and Europe. Our report extends its distribution range to Pakistan.

KEY WORDS—*Ascomycota*, edible fungi, *Morchellaceae*, phylogeny, systematics

Introduction

Malam Jabba is located in northern Pakistan between the Himalayan and Hindukush foothills in the Swat district of the Malakand Division in Khyber Pakhtunkhwa Province (KP). The area, with elevations ranging from 990 m at the valley entrance to 2880 m at the highest peak of Shagar Sar, is a mycologically rich region. In Pakistan, *Morchella* species fruit during spring at lower altitudes, but as late as July at higher elevations in snowfall regions (Hamayun & al. 2006, Badshah & al. 2015). True morels (*Morchella* spp.) belong to *Morchellaceae* Rchb. (*Pezizales*, *Ascomycota*) and are amongst the most highly valued edible fungi due to their excellent flavor and relatively short fruiting season (Taşkın & al. 2016). A high diversity of *Morchella* species has been described from various continents and regions, including Pakistan (Hamayun & al. 2006).

In recent years, a series of multilocus molecular phylogenetic studies of *Morchella* were performed employing phylogenetic species recognition based on genealogical concordance (GCPSR, sensu Taylor & al. 2000). These studies revealed considerable continental endemism and provincialism within the genus (O'Donnell & al. 2011; Du & al. 2012a,b; Richard & al. 2015; Loizides & al. 2016; Taşkın & al. 2010, 2012, 2016). This research also highlighted the difficulties surrounding morphology-based species identification and helped provide a robust framework for subsequent taxonomic revisions (Clowez 2012, 2014, 2015; Kuo & al. 2012; Richard & al. 2015; Voitek & al. 2016; Loizides & al. 2015, 2016). Multilocus molecular phylogenetic studies have so far revealed nearly 70 *Morchella* spp. worldwide (Loizides 2017).

A number of morel species—*Morchella conica*, *M. crassipes*, *M. elata*, *M. esculenta*, *M. rotunda*, *M. semilibera*—have been previously reported from the Kohistan and Swat regions of Pakistan based solely on morphology (Hamayun & al. 2006). These identifications based on old taxonomical concepts have not, however, been verified by molecular analysis. In this paper we report *Morchella pulchella* as a new record from Pakistan identified primarily using DNA sequence analysis.

Materials & methods

Specimens were collected during a field survey at Malam Jabba, District Swat, Province Khyber Pakhtunkhwa, Pakistan, in April 2015 under *Pinus wallichiana* near grasses. Tissues were mounted in glycerin or distilled water and examined microscopically using a Leitz HM Lux compound microscope under 400× magnification. The preserved specimens have been deposited in the herbarium of Plant Sciences, Quaid-i-Azam University, Islamabad, Pakistan (ISL).

DNA extraction and PCR amplification were conducted with the Sigma-Aldrich REDExtract-N-Amp™ Plant PCR Kit following the manufacturer's instructions. DNA was amplified with the following markers and primer pairs: translation elongation factor 1- α (TEF1) with EF526F & EF3AR (Rehner & Buckley 2005), RNA polymerase II large subunit (RPB1) with gRPB1A & aRPB1C (Matheny & al. 2002) and the RNA polymerase II second largest subunit (RPB2) with 9F & 3R (Liu & al. 1999). One μ L of genomic DNA was amplified in total volumes of 20 μ L in an Eppendorf Master Cycler Gradient thermocycler following the parameters of 94°C for 3 min, 35 cycles of 94°C for 30 s, 53°C for 30 s, 72°C for 1–2 min with a final

FIG. 1—Phylogenetic tree inferred from partial TEF1 sequences of *Morchella* spp., using the Neighbor Joining method. Posterior probabilities and bootstrap values are given on each node of the tree; the values for poorly supported nodes (bootstrap <40) are enclosed in parentheses. The new Pakistani sequence is set in bold font.



extension at 72°C for 7 min. Raw sequence data were edited using BioEdit (Hall 1999), assembled with Codon Code Aligner 4.1.1, and deposited in GenBank.

Sequences were analyzed using the BLASTn analysis tool (www.ncbi.nlm.nih.gov). Our newly generated *TEF1* sequence plus 26 *TEF1* sequences representing *Morchella* species previously deposited in NCBI were used for the phylogenetic analysis. DNA sequences were aligned using Muscle E multiple alignment tool within MEGA v. 7.0 (Kumar & al. 2016). A phylogenetic tree was inferred using the Neighbour-Joining method. Posterior probabilities were determined and the bootstrap analysis was based on 1000 pseudo-replicates using the NJ option.

Results

BLASTn analysis using the *TEF1* sequence (FIG. 1) as query matched three GenBank sequences representing *M. pulchella* (KM491180: 100% identity, 636/636 bp) and an unidentified *Morchella* sp. (GU551381: 100% identity, 636/636 bp; GU551390: 99% identity, 635/636 bp).

Similarly, the *RPB1* sequence matched three GenBank sequences representing *M. septentrionalis* (KM588048 & JX173414: 100% identity, 785/785 bp) and *M. pulchella* (HM056439: 100% identity, 785/785 bp).

Finally, the *RPB2* sequence matched three GenBank sequences representing *M. pulchella* (KM588041, 100% identity, 809/809 bp) and *M. septentrionalis* (KM588048 & JX173414: 100% identity, 805/805 bp).

Taxonomy

Morchella pulchella Clowez & Franç. Petit,

Bull. Soc. Mycol. Fr. 126: 314. 2012.

FIG. 2

ASCOMATA 60–70 × 52–56 mm, conical, brown. HYMENOPHORE 50 × 40 mm, slightly conical, ridged with 26 primary ridges per ascoma, ridges regular to irregular in shape, 2–5 mm long, 3 mm broad, pitted, pits (n = 55–60) short and elongated. Fertile area sharply attached to the stipe. STIPE hollow, white, convex, tapered at apex and wider at base with short rays. ASCOSPORES 20.4–24 × 13–15.6 µm, ellipsoid, smooth, with homogenous contents. ASCI 8-spored, cylindrical, hyaline, 200–260 × 15–20 µm, length/width (Qm) ratio 1.5 µm. PARAPHYSES shorter than asci, 78–150 × 10–14 µm, cylindrical, subclavate, apices rounded, narrow, mostly subacute, rarely acute, septate, hyaline with homogenous contents; yellow hyphal elements on sterile ridges 77–181 × 7.9–26 µm.

MATERIAL EXAMINED—PAKISTAN, KHYBER PAKHTUNKHWA, Swat District, Malam Jabba, 34°47'54"N 72°34'33"E, altitude 1800 m, under *Pinus wallichiana* A.B. Jacks. and local grasses, 15 April 2015, coll. Hussain Badshah B10 (ISL-67968; GenBank MF400855, MF400856, MF400857).




FIG. 2—*Morchella pulchella* (ISL-67968)

A: ascocarp. B, D: asci with ascospores. C: paraphyses.

Discussion

The phylogenetic analysis (FIG. 1) clustered sequences from our collection with *M. pulchella*, supporting the Malam Jabba specimen as a new record of that species from Pakistan. As in previous studies (Du & al. 2012, Taşkın & al. 2012, Richard & al. 2015), our phylogenetic analysis provides no support for the reciprocal monophyly of *M. pulchella* and *M. septentrionalis* and emphasizes the need to investigate further the taxonomic status of *M. pulchella* and *M. septentrionalis*.

Morchella pulchella has previously been reported only from China (Du & al. 2012a,b), Turkey (Taşkın & al. 2010), France (Clowez 2012, Richard

& al. 2015), and Spain (Alonso & al. 2016). Confirmation of *M. pulchella* from an apparently undisturbed mountainous site in Pakistan suggests that *M. pulchella* is unlikely to have been introduced to this region through human activity and has an innately Eurasian distribution.

Acknowledgements

The authors are sincerely thankful to Dr. Kerry O'Donnell (NCAUR, USDA Agricultural Research Service, Peoria, IL) and Dr. Xi-Hui Du (Chongqing Normal University, China) for acting as presubmission reviewers and giving valuable suggestions. We would also like to acknowledge Prof. Donald H. Pfister (Farlow Library and Herbarium, Harvard University) for linguistic improvement and Dr. Michael Loizides (Cyprus Mycological Association, Nicosia) and Dr. Shaun Pennycook (Manaaki Whenua Landcare Research, Auckland, New Zealand) for valuable suggestions on the manuscript.

Literature cited

- Alonso LJ, Fernández A, Monedero C. 2017. *Morchella pulchella*, una especie poco frecuente. YESCA Revista de micología 28: 47–52.
- Badshah H, Ullah F, Khan MU, Mumtaz AS, Malik RN. 2015. Pharmacological activities of selected wild mushrooms in South Waziristan (FATA), Pakistan. South African Journal of Botany 97: 107–110. <https://doi.org/10.1016/j.sajb.2014.12.002>
- Clowez P. 2012. Les morilles: une nouvelle approche mondiale du genre *Morchella*. Bulletin Trimestriel de la Société Mycologique de France 126(3–4): 199–376.
- Clowez P, Alvarado P, Becerra M, Bilbao T, Moreau PA. 2014. *Morchella fluvialis* sp. nov. (*Ascomycota*, *Pezizales*): a new but widespread morel in Spain. Bulletin de Sociedad Micologica de Madrid 38: 251–260.
- Clowez P, Bellanger JM, de la Osa LR, Moreau PA. 2015. *Morchella palazonii* sp. nov. (*Ascomycota*, *Pezizales*): une nouvelle morille méditerranéenne. Clé des *Morchella* sect. *Morchella* en Europe. Documents Mycologiques 36: 71–84.
- Du X-H, Zhao Q, O'Donnell K, Rooney AP, Yang ZL. 2012a. Multigene molecular phylogenetics reveals true morels (*Morchella*) are especially species-rich in China. Fungal Genetics and Biology 49: 455–469. <https://doi.org/10.1016/j.fgb.2012.03.006>
- Du X-H, Zhao Q, Yang ZL, Hansen K, Taşkın H, Büyükalaca S, Dewsbury D, Moncalvo JM, Douhan GW, Robert VARG, Crous PW, Rehner SA, Rooney AP, Sink S, O'Donnell K. 2012b. How well do ITS rDNA sequences differentiate species of true morels (*Morchella*)? Mycologia 104(6): 1351–1368. <https://doi.org/10.3852/12-056>
- Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic Acids Symposium Series 41: 95–98. [London]: Information Retrieval Ltd., c1979–c2000.
- Hamayun M, Khan SA, Sohn EY, Lee IJ. 2006. Folk medicinal knowledge and conservation status of some economically valued medicinal plants of district Swat, Pakistan. Lyonia 11: 101–113.
- Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. Molecular Biology and Evolution 33: 1870–1874. <https://doi.org/10.1093/molbev/msw054>

- Kuo M, Dewsbury DR, O'Donnell K, Carter MC, Rehner SA, Moore JD, Moncalvo JM, Canfield SA, Stephenson SL, Methven AS, Volk TJ. 2011. Taxonomic revision of true morels (*Morchella*) in Canada and the United States. *Mycologia* 104: 1159–1177. <https://doi.org/10.3852/11-375>
- Liu YJ, Whelen S, Hall BD. 1999. Phylogenetic relationships among ascomycetes: evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16(12): 1799–1808. <https://doi.org/10.1093/oxfordjournal.molbev.a026092>
- Loizides M. 2017. Morels: the story so far. *Field Mycology* 18(2): 42–53. <https://doi.org/10.1016/j.fldmyc.2017.04.004>
- Loizides M, Alvarado P, Clowez P, Moreau PA, de la Osa LR, Palazón A. 2015. *Morchella tridentina*, *M. rufobrunnea*, and *M. kakiicolor*: a study of three poorly known Mediterranean morels, with nomenclatural updates in section *Distantes*. *Mycological Progress* 14:13 [18 p.]. <https://doi.org/10.1007/s11557-015-1030-6>
- Loizides M, Bellanger JM, Clowez P, Richard P, Moreau PA. 2016. Combined phylogenetic and morphological studies of true morels (*Pezizales*, *Ascomycota*) in Cyprus reveal significant diversity, including *Morchella arbutiphila* and *M. disparilis* spp. nov. *Mycological Progress* 15:39 [28 p.]. <https://doi.org/10.1007/s11557-016-1180-1>
- Matheny PB, Liu YJ, Ammirati JF, Hall BD. 2002. Using *RPB1* sequences to improve phylogenetic inference among mushrooms (*Inocybe*, *Agaricales*). *American Journal of Botany* 89: 688–698. <https://doi.org/10.3732/ajb.89.4.688>
- O'Donnell K, Rooney AJ, Mills GL, Kuo M, Weber NS, Rehner SA. 2011. Phylogeny and historical biogeography of true morels (*Morchella*) reveals an early Cretaceous origin and high continental endemism and provincialism in the Holarctic. *Fungal Genetics and Biology* 48: 252–265. <https://doi.org/10.1016/j.fgb.2010.09.006>
- Rehner SA, Buckley E. 2005. A *Beauveria* phylogeny inferred from nuclear ITS and EF1- α sequences: evidence for cryptic diversification and links to *Cordyceps* teleomorphs. *Mycologia* 97: 84–98. <https://doi.org/10.1080/15572536.2006.11832842>
- Richard F, Bellanger JM, Clowez P, Hansen K, O'Donnell K, Urban A, Sauve M, Courtecuisse R, Moreau PA. 2015. True morels (*Morchella*, *Pezizales*) of Europe and North America: evolutionary relationships inferred from multilocus data and a unified taxonomy. *Mycologia* 107: 359–382. <https://doi.org/10.3852/14-166>
- Taşkın H, Büyükalaca S, Doğan HH, Rehner SA, O'Donnell K. 2010. A multigene molecular phylogenetic assessment of true morels (*Morchella*) in Turkey. *Fungal Genetics and Biology* 47: 672–682. <https://doi.org/10.1016/j.fgb.2010.05.004>
- Taşkın H, Büyükalaca S, Hansen K, O'Donnell K. 2012. Multilocus phylogenetic analysis of true morels (*Morchella*) reveals high levels of endemics in Turkey relative to other regions of Europe. *Mycologia* 104: 446–461. <https://doi.org/10.3852/11-180>
- Taşkın H, Doğan HH, Büyükalaca S, Clowez P, Moreau PA, O'Donnell K. 2016. Four new morel (*Morchella*) species in the elata subclade (*M.* sect. *Distantes*) from Turkey. *Mycotaxon* 131: 467–482. <https://doi.org/10.5248/131.467>
- Taylor JW, Jacobsen DJ, Kroken S, Kasuga T, Geiser DM, Hibbett DS, Fisher MC. 2000. Phylogenetic species recognition and species concepts in fungi. *Fungal Genetics and Biology* 31: 21–32.
- Voitk A, Beug MW, O'Donnell K, Burzynski M. 2016. Two new species of true morels from Newfoundland and Labrador: cosmopolitan *Morchella eohespera* and parochial *M. laurentiana*. *Mycologia* 108(1): 31–37. <https://doi.org/10.3852/15-149>